

Since publication of her article, the author reports no further potential conflict of interest.

1. Robinson JC. Price transparency begins at home. *Front Health Serv Manage* 2007;23:25-8.
2. O'Connor AM, Bennett CL, Stacey D, et al. Decision aids for people facing health treatment or screening decisions. *Cochrane Database Syst Rev* 2009;3:CD001431.
3. Gellad ZF, Lyles KW. Direct-to-consumer advertising of pharmaceuticals. *Am J Med* 2007;120:475-80.

4. Hermer LD, Brody H. Defensive medicine, cost containment, and reform. *J Gen Intern Med* 2010;25:470-3.
5. Grumbach K, Bodenheimer T, Grundy P. The outcomes of implementing patient-centered medical home interventions: a review of the evidence on quality, access and costs from recent prospective evaluation trials, August 2009. Washington, DC: Patient-Centered Primary Care Collaborative, 2009. (Accessed August 5, 2010, at http://www.pcpcc.net/files/pcmh_evidence_outcomes_2009.pdf)

Reference Range for Cerebrospinal Fluid Opening Pressure in Children

TO THE EDITOR: A reference range for cerebrospinal fluid (CSF) opening pressure in children undergoing diagnostic lumbar puncture has not been established.¹ The influence of age, body-mass index (BMI), and depth of sedation on opening pressure in children is also uncertain.²

We conducted a 2-year, single-center prospective study of CSF opening pressure in children undergoing lumbar puncture as part of their routine clinical care. Our objective was to establish a useful, clinically relevant reference range for opening pressure and a threshold value for abnormally elevated opening pressure (in the 90th percentile or higher, defined a priori). Our reference group consisted of children between 1 and 18 years of age, none of whom was taking a medication or had signs of a disease or condition that would alter opening pressure on lumbar puncture (e.g., papilledema, hydrocephalus, cerebral edema, Chiari malformation, meningitis, or use of diuretics). The procedure was performed with subjects in the lateral recumbent position, and the opening pressure was measured with the use of a standard manometer. (For enrollment criteria, methods, and results, see the Supplementary Appendix, available with the full text of this letter at [NEJM.org](http://www.nejm.org).) Reference-range parameters for opening pressure were determined with the use of standard descriptive statistics, whereas associations between age, BMI, and depth of sedation were determined using linear regression analysis.

We enrolled 472 subjects during the study period, and 197 (41.7%) met the inclusion criteria for the reference-range population. The threshold for an abnormally elevated opening pressure, deter-

mined on the basis of the 90th percentile for all patients in the reference population, was 28 cm of water (Fig. 1A). The threshold for an abnormally reduced pressure in the 10th percentile was 11.5 cm of water. Subjects placed under moderate to deep sedation during lumbar puncture had a slightly higher opening pressure as compared with those not receiving any sedative medication ($\beta=3.459$, $P=0.002$) (Fig. 1B). Similar to studies in adults,^{3,4} our study showed a small yet statistically significant positive relationship between opening pressure and BMI ($\beta=0.313$, $P=0.002$) (Fig. 1C). Subject age was not associated with increased opening pressure ($P=0.43$) (Fig. 1D). Given the significant association between depth of sedation and BMI on opening pressure, a post-hoc analysis of opening pressure percentiles was calculated for the 52 subjects who received minimal or no sedation and were not classified as obese, resulting in a 90th percentile of 25 cm of water.

This study systematically defined the reference range for CSF opening pressure and the threshold value for abnormally elevated opening pressure in a representative pediatric population (in whom evident pathology was excluded) undergoing diagnostic lumbar puncture. It is our opinion that previous estimates of what constitutes an "abnormal" opening pressure, such as 20 cm of water,^{1,5} should be reconsidered, and that for most children an opening pressure above 28 cm of water should be considered elevated.

Robert A. Avery, D.O.

Samir S. Shah, M.D., M.S.C.E.

Daniel J. Licht, M.D.

Jeffrey A. Seiden, M.D.




Figure 1. Distribution of CSF Opening Pressure and Effects of Sedation, Body-Mass Index, and Age.

Panel A shows the distribution of the cerebrospinal fluid (CSF) opening pressure for 197 study subjects who met the inclusion criteria for the reference group. The mean (\pm SD) CSF opening pressure was 19.8 ± 6.8 cm of water. The values ranged from 6 to 47 cm of water and were similar in subjects 10 years of age or older and those younger than 10 years. The threshold for an abnormally elevated opening pressure, which was based on subjects with opening pressures in the 90th percentile or higher, was 28 cm of water. The threshold for an abnormally low opening pressure, based on subjects in the 10th percentile, was 11.5 cm of water. Panel B shows the results of an unadjusted linear regression analysis of the relationship between opening pressure and sedation ($\beta=3.459$, $P=0.005$). The relationship between opening pressure and body-mass index (BMI, the weight in kilograms divided by the square of the height in meters) is shown in Panel C ($\beta=0.138$, $P=0.12$) and that between opening pressure and age in Panel D ($\beta=0.328$, $P<0.001$). Statistically significant associations between opening pressure and moderate-to-deep sedation ($\beta=4.990$, $P=0.002$) and opening pressure and BMI ($\beta=0.313$, $P=0.002$) remained in multivariable analysis adjusted for age. In a multivariable linear regression model, no significant association was found between age ($\beta=0.097$, $P=0.42$) and opening pressure, even when age was categorized as 10 years or more or as less than 10 years ($\beta=1.141$, $P=0.30$). No significant association was found between opening pressure and sedation with ketamine ($\beta=3.080$; 95% confidence interval, -0.551 to 6.712 ; $P=0.10$).

Jimmy W. Huh, M.D.
Jan Boswinkel, M.D.
Michael D. Ruppe, M.D.
Amber Chew, M.P.H.
Rakesh D. Mistry, M.D.
Grant T. Liu, M.D.
Children's Hospital of Philadelphia
Philadelphia, PA
ravery@cnmc.org

The content of this letter is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Disclosure forms provided by the authors are available with the full text of this article at NEJM.org.

Supported by funds from the National Institutes of Health—National Eye Institute loan repayment program and the 1T32NS061779-01 Neurologic Clinical Epidemiology Training Program (to Dr. Avery); a grant from the National Institute of Allergy and Infectious Diseases (K01 AI73729) and the Robert Wood Johnson Foundation, under its Physician Faculty Scholar Program (to Dr. Shah); grants from the National Institute of Neurological Disorders and Stroke (NINDS) (NS-052380) and the Dana Foundation and a gift from the June and Steve Wolfson Fund for Neurological Research (to Dr. Licht); and a grant from NINDS (NS-053651) (to Dr. Huh).

1. Rangwala LM, Liu GT. Pediatric idiopathic intracranial hypertension. *Surv Ophthalmol* 2007;52:597-617.
2. Friedman DI, Jacobson DM. Idiopathic intracranial hypertension. *J Neuroophthalmol* 2004;24:138-45.
3. Corbett JJ, Mehta MP. Cerebrospinal fluid pressure in normal obese subjects and patients with pseudotumor cerebri. *Neurology* 1983;33:1386-8.
4. Whiteley W, Al-Shahi R, Warlow CP, Zeidler M, Lueck CJ. CSF opening pressure: reference interval and the effect of body mass index. *Neurology* 2006;67:1690-1.
5. Nechyba C. Blood chemistries and body fluids. In: Gunn VL, Nechyba C, eds. *The Harriet Lane handbook: a manual for pediatric house officers*. 16th ed. Philadelphia: Mosby, 2002:549-59.

Correspondence Copyright © 2010 Massachusetts Medical Society.

INSTRUCTIONS FOR LETTERS TO THE EDITOR

Letters to the Editor are considered for publication, subject to editing and abridgment, provided they do not contain material that has been submitted or published elsewhere. Please note the following:

- Letters in reference to a *Journal* article must not exceed 175 words (excluding references) and must be received within 3 weeks after publication of the article.
- Letters not related to a *Journal* article must not exceed 400 words.
- A letter can have no more than five references and one figure or table.
- A letter can be signed by no more than three authors.
- Financial associations or other possible conflicts of interest must be disclosed. Disclosures will be published with the letters. (For authors of *Journal* articles who are responding to letters, we will only publish new relevant relationships that have developed since publication of the article.)
- Include your full mailing address, telephone number, fax number, and e-mail address with your letter.
- All letters must be submitted at authors.NEJM.org.

Letters that do not adhere to these instructions will not be considered. We will notify you when we have made a decision about possible publication. Letters regarding a recent *Journal* article may be shared with the authors of that article. We are unable to provide prepublication proofs. Submission of a letter constitutes permission for the Massachusetts Medical Society, its licensees, and its assignees to use it in the *Journal's* various print and electronic publications and in collections, revisions, and any other form or medium.

CORRECTIONS

Acupuncture for Chronic Low Back Pain (July 29, 2010;363:454-61). The middle initial of the second author (page 454) was incorrect as published; the correct name is Helene M. Langetvin. The article has been corrected at NEJM.org.

A Randomized Trial of Treatment for Acute Anterior Cruciate Ligament Tears (July 22, 2010;363:331-42). In Figure 1 (page 335), the last line in the box just below "69 Were assigned to undergo rehabilitation and early ACL reconstruction" should have read, "1 Had intact ACL," rather than "2 Had intact ACL." We regret the error. The article has been corrected at NEJM.org.

Early-Stage Hodgkin's Lymphoma (August 12, 2010;363:653-62). In the print edition of the *Journal*, there were a number of instances of "therapy" that should have been "radiotherapy." We regret the error. The article was corrected before publication at NEJM.org; subscribers to the print edition of the *Journal* will receive a corrected reprint.

NOTICES

Notices submitted for publication should contain a mailing address and telephone number of a contact person or department. We regret that we are unable to publish all notices received. Notices also appear on the Journal's Web site (NEJM.org/medical-conference). The listings can be viewed in their entirety or filtered by specialty, location, or month.

CALL FOR APPLICATIONS

The American Philosophical Society is accepting applications for its "Daland Fellowships in Clinical Investigation," which support research in the several branches of clinical medicine, including internal medicine, neurology, pediatrics, psychiatry, and surgery.

Contact Linda Musumeci, Daland Fellowships in Clinical Investigation, American Philosophical Society, 104 S. Fifth St., Philadelphia, PA 19106-3386; or call (215) 440-3429; or e-mail lmusumeci@amphilsoc.org; or see <http://www.amphilsoc.org/grants/daland>.

MAYO CLINIC SCOTTSDALE

The following courses will be offered in Sedona, AZ: "Mayo Clinic 13th Annual Internal Medicine Update: Sedona 2010" (Oct. 7-10 or Oct. 21-24) and "7th Annual Women's Health Update" (March 10-12).

Contact Mayo Clinic Scottsdale, 13400 E. Shea Blvd., Scottsdale, AZ 85259; or call (480) 301-4580.

CALL FOR APPLICATIONS

The American Society of Tropical Medicine and Hygiene is accepting applications for its "Postdoctoral Fellowships in Tropical Infectious Diseases," which provide support for individuals to conduct research in tropical infectious diseases or other clinical conditions unique to tropical medicine. Deadline for submission is Sept. 1.

Contact the American Society of Tropical Medicine and Hygiene, 111 Deer Lake Rd., Suite 100, Deerfield, IL 60015; or call (847) 480-9592; or fax (847) 480-9282; or e-mail info@astmh.org; or see <http://www.astmh.org>.

39TH ANNUAL MEETING OF THE AMERICAN COLLEGE OF CLINICAL PHARMACOLOGY

The meeting will be held in Baltimore, Sept. 12-14.

Contact Mina Behari or Krista Levy, American College of Clinical Pharmacology, 3 Ellinwood Court, New Hartford, NY 13413-1105; or call (315) 768-6117; or fax (315) 768-6119; or e-mail mina@accp1.org; or see <http://www.accp1.org>.

HARVARD LUNG CONFERENCE

The conference will be held in Boston, Oct. 6 and 7, in conjunction with the "CIMIT Inhalation Technology Workshop" on Oct. 8.

Contact Tia Radha Dastidar, Strategic Results, 3225 Ellerslie Ave., Suite 303E, Baltimore, MD 21218; or call (443) 451-7252; or e-mail tiaradha@strategicresults.com; or see <http://www.harvardlungconference.org>.

OPTIMIZING MECHANICAL VENTILATION 2010: A HANDS-ON PRACTICAL EMPHASIS

The conference will be held in St. Paul, MN, Oct. 8-10.

Contact Sharon Kopp-Huth, HealthPartners Institute for Medical Education, Center for Continuing Professional Development, P.O. Box 1309, MC21101B, Minneapolis, MN 55440-1309; or call (952) 883-6224; or e-mail sharon.a.kopp@healthpartners.com; or see <http://MechanicalVentilationConferenceMN.com>.